

Finite set of invariants to characterize local Clifford equivalence of stabilizer states

Maarten Van den Nest,* Jeroen Dehaene, and Bart De Moor
Katholieke Universiteit Leuven, ESAT-SCD, Belgium.

(Dated: February 9, 2008)

The classification of stabilizer states under local Clifford (LC) equivalence is of particular importance in quantum error-correction and measurement-based quantum computation. Two stabilizer states are called LC equivalent if there exists a local Clifford operation which maps the first state to the second. We present a finite set of invariants which completely characterizes the LC equivalence class of any stabilizer state. Our invariants have simple descriptions within the binary framework in which stabilizer states are usually described.

PACS numbers: 03.67.-a

Stabilizer states constitute a class of multipartite pure quantum states which is of considerable interest in quantum information theory (QIT) and quantum computing (QC). A stabilizer state on n qubits is defined as a simultaneous eigenvector of a maximal set of commuting observables in the Pauli group, where the latter is the group generated by all n -fold tensor products of the Pauli matrices and the identity. To name but two of their presently most prominent applications, stabilizer states firstly appear as codewords of quantum stabilizer codes in the theory of quantum error-correction [1], and, secondly, they are the resources for QC in a computation model where only single-qubit measurements are performed (the one-way quantum computer [2]). In order to understand the role of stabilizer states in these and other QIT and QC tasks, the mathematical and physical properties of these states have recently been studied in a number of accounts [3, 4, 5, 6, 7, 8]. One central problem in this research is the classification of stabilizer states under local unitary (LU) equivalence. Indeed, next to the natural relevance of this issue in the study of the entanglement properties of stabilizer states, it is also of particular importance both in the development of the one-way quantum computer and in the coding theoretic aspect of stabilizer states. In this paper we consider a restricted version of LU equivalence of stabilizer states, in that we consider only those local unitary operations which belong to the local Clifford group (LC). The local Clifford group consists of all local unitary operators which map the Pauli group to itself under conjugation. We will call two stabilizer states LC equivalent if there exists a local Clifford operator which maps the first state to the second.

The restriction of considering LC equivalence rather than general LU equivalence has two main motivations. Firstly, the problem of general LU equivalence is difficult and it is therefore appropriate to first consider a more manageable subproblem, for which LC equivalence is a suitable candidate - given the very explicit connections between stabilizer states, the Pauli group and the (local) Clifford group. Secondly, this so-called 'subproblem'

might in fact not be a subproblem at all, as the natural question exists whether every two LU equivalent stabilizer states are necessarily LC equivalent. Although this question is to date unanswered in general, our work [9], following work performed in Ref. [10], shows that this assertion is at least true for some interesting subclasses of stabilizer states and a positive answer to this question is widely believed.

In the following we study LC invariants of stabilizer states, i.e. functions in the entries of a stabilizer state which take on equal values in LC equivalent states. We construct a finite *complete set* of invariants, which characterizes the LC equivalence class of any stabilizer state. Such a study of LC invariants primarily serves to obtain a characterization of LC equivalence classes of stabilizer states and to gain insight in the structure of these classes, and to a lesser extent to obtain algorithms which can recognize LC equivalence of two given states. Indeed, an efficient algorithm to recognize LC equivalence of stabilizer states has recently been presented in our earlier work and there is little hope that equally efficient algorithms can be constructed on the basis of invariants (see below). Therefore, we believe that the main merit of the present result is, firstly, that it gives a *finite* characterization of the LC equivalence class of any stabilizer state and, secondly, that our invariants exhibit a very transparent structure in terms of the stabilizer formalism.

Let us now start by recalling the basic notions concerning stabilizer states and the local Clifford group. A stabilizer state $|\psi\rangle$ on n qubits is the unique simultaneous eigenvector with eigenvalue 1 of a set of n commuting and independent observables in the Pauli group $\mathcal{G}_n$. The latter consists of all operators of the form $\alpha\sigma_{a_1} \otimes \cdots \otimes \sigma_{a_n}$, where $\alpha \in \{\pm 1, \pm i\}$ is an overall phase factor and σ_{a_i} is either the 2×2 identity matrix or one of the Pauli matrices $\sigma_x, \sigma_y, \sigma_z$, for every $i = 1, \dots, n$. The stabilizer $\mathcal{S}(\psi)$ of $|\psi\rangle$ is the set of all Pauli operators which have $|\psi\rangle$ as an eigenvector with eigenvalue 1. It is well known (see e.g. [11]) that the stabilizer formalism has an equivalent formulation in terms of algebra over the field $\mathbb{F}_2 = \text{GF}(2)$, where arithmetic is performed modulo 2. The heart of this binary representation is an encoding of the Pauli matrices by pairs of bits, writing $\sigma_0 = \sigma_{00}$,

*Electronic address: maarten.vandennest@esat.kuleuven.ac.be

$\sigma_x = \sigma_{01}$, $\sigma_z = \sigma_{10}$ and $\sigma_y = \sigma_{11}$ and subsequently

$$\sigma_{w_1 w'_1} \otimes \cdots \otimes \sigma_{w_n w'_n} =: \sigma_{(w, w')}, \quad (1)$$

where $w := (w_1, \dots, w_n)$, $w' := (w'_1, \dots, w'_n) \in \mathbb{F}_2^n$ are n -dimensional binary vectors. Note that the information about the i th qubit is distributed over the i th components of the vectors w and w' . Identifying every Pauli operator in an n -qubit stabilizer $\mathcal{S}(\psi)$ with its $2n$ -dimensional binary index vector, one can show that $\mathcal{S}(\psi)$ corresponds to an n -dimensional self-dual linear subspace S_ψ of $\mathbb{F}_2^{2n}$; the self-duality of this subspace is with respect to a symplectic inner product on $\mathbb{F}_2^{2n}$.

The Clifford group $\mathcal{C}_1$ on one qubit is the group of all 2×2 unitary operators which map σ_u to $\alpha_u \sigma_{\pi(u)}$ under conjugation, where $u = x, y, z$, for some $\alpha_u = \pm 1$ and some permutation π of $\{x, y, z\}$. The local Clifford group $\mathcal{C}_n^l$ on n qubits is the n -fold tensor product of $\mathcal{C}_1$ with itself. In the binary stabilizer framework, local Clifford operations $U \in \mathcal{C}_n^l$ correspond to nonsingular $2n \times 2n$ binary matrices Q of the block form

$$Q = \begin{bmatrix} A & B \\ C & D \end{bmatrix}, \quad (2)$$

where the $n \times n$ blocks A, B, C, D are diagonal [5]. We denote the diagonal entries of A, B, C, D by a_i, b_i, c_i, d_i , respectively. The n submatrices

$$Q_i := \begin{bmatrix} a_i & b_i \\ c_i & d_i \end{bmatrix} \in GL(2, \mathbb{F}_2) \quad (3)$$

correspond to the tensor factors of U . We denote the group of all such Q by $\mathcal{C}_n^l$. In the binary stabilizer framework, two n -qubit stabilizer states $|\psi\rangle, |\psi'\rangle$ are LC equivalent if and only if there exists an operator $Q \in \mathcal{C}_n^l$ such that $Q S_\psi = S_{\psi'}$, i.e., Q maps the space S_ψ to the space $S_{\psi'}$.

Finally, let us recall the definition of *support* in this context of stabilizers, which will be of considerable importance in the following. The support $\text{supp}(v)$ of any vector $v \in \mathbb{F}_2^{2n}$ is the set

$$\text{supp}(v) = \{i \in \{1, \dots, n\} \mid (v_i, v_{n+i}) \neq (0, 0)\}.$$

Note that $\text{supp}(v)$ contains exactly those $i \in \{1, \dots, n\}$ such that the i th tensor factor of σ_v differs from the identity.

We are now in a position to state the first result of this paper. In theorem 1 a finite set of invariants which characterizes the LC equivalence class of any stabilizer state, is presented.

Theorem 1 *Let $|\psi\rangle$ be a stabilizer state on n qubits. Let $r \in \mathbb{N}_0$ and consider subsets $\omega^k, \omega^{kl} \subseteq \{1, \dots, n\}$ for every $k, l \in \{1, \dots, r\}$ with $k < l$. Denote $\Omega := (\omega^1, \omega^2, \dots, \omega^{12}, \omega^{13}, \dots)$ and let $\mathcal{T}_{n,r}^\Omega(\psi)$ be the set consisting of all tuples $(v^1, \dots, v^r) \in S_\psi \times \cdots \times S_\psi$ satisfying*

$$\text{supp}(v^k) = \omega^k, \quad \text{supp}(v^k + v^l) = \omega^{kl}. \quad (4)$$

Then (i) $|\mathcal{T}_{n,r}^\Omega(\psi)|$ is an LC invariant and (ii) the LC equivalence class of $|\psi\rangle$ is completely determined by the values of all invariants $|\mathcal{T}_{n,n}^\Omega(\psi)|$ (i.e. where $r = n$).

Proof: Statement (i) is trivial. To prove (ii), let $|\psi\rangle$ and $|\psi'\rangle$ be two stabilizer states on n qubits such that $|\mathcal{T}_{n,n}^\Omega(\psi)| = |\mathcal{T}_{n,n}^\Omega(\psi')|$ for every tuple Ω as defined above. We will show that this implies that $|\psi\rangle$ and $|\psi'\rangle$ are LC equivalent. Fix a basis $\{v^1, \dots, v^n\}$ of S_ψ and define the tuple of sets $\Omega_* = (\omega_*^1, \dots, \omega_*^{12}, \dots)$ such that

$$\omega_*^k = \text{supp}(v^k), \quad \omega_*^{kl} = \text{supp}(v^k + v^l), \quad (5)$$

for every $k, l = 1, \dots, n$ with $k < l$. It follows that $|\mathcal{T}_{n,n}^{\Omega_*}(\psi)|$ is nonzero, and therefore $|\mathcal{T}_{n,n}^{\Omega_*}(\psi')|$ is nonzero as well, since these numbers are equal by assumption. Therefore, there exist vectors $w^1, \dots, w^n \in S_{\psi'}$ such that

$$\text{supp}(w^k) = \omega_*^k, \quad \text{supp}(w^k + w^l) = \omega_*^{kl}, \quad (6)$$

for every k, l . Expressions (5) and (6) show that

$$\begin{aligned} (v_i^k, v_{n+i}^k) &= (0, 0) \quad \text{iff} \quad (w_i^k, w_{n+i}^k) = (0, 0) \\ (v_i^k, v_{n+i}^l) &= (v_i^l, v_{n+i}^l) \quad \text{iff} \quad (w_i^k, w_{n+i}^k) = (w_i^l, w_{n+i}^l), \end{aligned}$$

for every $i, k, l = 1, \dots, n$. This implies the existence of n operators $Q_i \in GL(2, \mathbb{F}_2)$ such that

$$Q_i \begin{bmatrix} v_i^k \\ v_{n+i}^k \end{bmatrix} = \begin{bmatrix} w_i^k \\ w_{n+i}^k \end{bmatrix} \quad (7)$$

for every $i, k = 1, \dots, n$, and the operators Q_i consequently constitute an operator $Q \in \mathcal{C}_n^l$ such that $Q v^k = w^k$ for every $k = 1, \dots, n$. Therefore, the space $Q S_\psi$ is spanned by the vectors $w^1, \dots, w^n$. Moreover, these vectors are linearly independent, as they are the images of a linearly independent set under an invertible linear transformation. Therefore, $\{w^1, \dots, w^n\}$ is a basis of $S_{\psi'}$ (since the latter is an n -dimensional vector space) and we obtain $Q S_\psi = S_{\psi'}$. This proves the theorem. $\square$

Let us briefly discuss this result. The invariants in theorem 1 are presented as cardinalities of certain subsets of $S_\psi \times \cdots \times S_\psi$, which are defined in terms of simple constraints on the supports of their elements. Note that for $r = 1$ these invariants simply count the number of elements in the stabilizer with a prescribed support, i.e., for every $\omega \subseteq \{1, \dots, n\}$ one has an invariant

$$|\{v \in S_\psi \mid \text{supp}(v) = \omega\}|. \quad (8)$$

These invariants are in fact 'local versions' of the so-called *weight distribution* of a stabilizer, a well known concept in quantum (and classical) coding theory which is e.g. used to produce bounds on how good codes can be (see e.g. [12]). Our invariants for $r \geq 2$ are in this respect generalizations of this well known notion.

While the structure of the invariants in theorem 1 is indeed very transparent, a direct calculation of the numbers $|\mathcal{T}_{n,r}^\Omega(\psi)|$ is likely to be hard. It is in this context interesting to consider the following variant of theorem 1, which presents an alternative complete family of LC invariants:

Theorem 2 *Using the same notations as in theorem 1, let $\mathcal{V}_{n,r}^\Omega(\psi)$ be the vector space consisting of all tuples $(v^1, \dots, v^r) \in S_\psi \times \dots \times S_\psi$ satisfying*

$$\text{supp}(v^k) \subseteq \omega^k, \quad \text{supp}(v^k + v^l) \subseteq \omega^{kl}. \quad (9)$$

Then (i) the dimension of $\mathcal{V}_{n,r}^\Omega(\psi)$ is an LC invariant and (ii) the LC equivalence class of $|\psi\rangle$ is completely determined by the values of all invariants $\dim(\mathcal{V}_{n,n}^\Omega(\psi))$.

Proof: Statement (i) is again trivial. We now prove (ii). Let $\omega \subseteq \{1, \dots, n\}$ and define the functions $\delta_\omega, \epsilon_\omega : \mathbb{F}_2^{2n} \rightarrow \mathbb{C}$, respectively, by $\delta_\omega(w) = 1$ if $\text{supp}(w) = \omega$ and $\delta_\omega(w) = 0$ otherwise, and $\epsilon_\omega(w) = 1$ if $\text{supp}(w) \subseteq \omega$ and $\epsilon_\omega(w) = 0$ otherwise. It is then straightforward to show the following relations:

$$\begin{aligned} \epsilon_\omega &= \sum_{\omega' \subseteq \omega} \delta_{\omega'} \\ \delta_\omega &= (-1)^{|\omega|} \sum_{\omega' \subseteq \omega} (-1)^{|\omega'|} \epsilon_{\omega'}, \end{aligned} \quad (10)$$

the first of which is trivial and the second of which can be verified using straightforward combinatorics. Letting Ω be a tuple of sets $\omega^k, \omega^{kl} \subseteq \{1, \dots, n\}$, with $k, l = 1, \dots, r$ and $k < l$, it follows that

$$\begin{aligned} |\mathcal{T}_{n,r}^\Omega(\psi)| &= \sum \left\{ \prod_k \delta_{\omega^k}(v^k) \right\} \left\{ \prod_{k < l} \delta_{\omega^{kl}}(v^k + v^l) \right\} \\ |\mathcal{V}_{n,r}^\Omega(\psi)| &= \sum \left\{ \prod_k \epsilon_{\omega^k}(v^k) \right\} \left\{ \prod_{k < l} \epsilon_{\omega^{kl}}(v^k + v^l) \right\}, \end{aligned}$$

where the sums run over all r -tuples $(v^1, \dots, v^r) \in S_\psi^{\times r}$. Using (10) one finds that every invariant $|\mathcal{T}_{n,r}^{\Omega_0}(\cdot)|$ is a linear combination of the invariants $\{|\mathcal{V}_{n,r}^\Omega(\cdot)|\}_\Omega$ with constant coefficients (and vice versa). The proof then readily follows after applying theorem 1 and noting that the dimension of any vector space $\mathcal{V}$ over $\mathbb{F}_2$ is given by $\dim(\mathcal{V}) = \log_2 |\mathcal{V}|$. $\square$

The main advantage of the invariants $\dim(\mathcal{V}_{n,r}^\Omega(\psi))$ over the $|\mathcal{T}_{n,r}^\Omega(\psi)|$'s is that in the former case one is dealing with (dimensions of) vector spaces, which are much more manageable from a computational point of view. Let us illustrate this with a simple example: take $\omega \subseteq \{1, \dots, n\}$ as before and consider the invariant

$$|\{v \in S_\psi \mid \text{supp}(v) \subseteq \omega\}|, \quad (11)$$

which is the counterpart of (8) in theorem 2. This invariant can indeed be calculated efficiently as follows: let $v^1, \dots, v^n$ be a basis of S_ψ and let $S = [v^1 \dots v^n]$ be the $2n \times n$ binary matrix which has the vectors v^j as its columns (the matrix S is generally referred to as a *generator matrix* of the stabilizer). Denoting by S_i^T the $2 \times n$ submatrix of S which is obtained by assembling the i th and the $(n+i)$ th row of S , for every $i = 1, \dots, n$, one readily verifies that the invariant (11) is equal to the corank

(i.e. the dimension of the kernel) over $\mathbb{F}_2$ of the matrix $S_\omega := (S_j^T)_{j \in \bar{\omega}}$, where $\bar{\omega}$ denotes the complement of ω in $\{1, \dots, n\}$. Note that the calculation of the (co)rank of a matrix over any field indeed requires only a polynomial number of operations in the dimensions of this matrix. This shows that the invariant (11) can easily be calculated when a generator matrix of the stabilizer is known (note that a stabilizer state is indeed usually presented in terms of a generator matrix). A straightforward continuation of the above argument shows that, e.g. for $r = 2$, an invariant $\dim(\mathcal{V}_{n,2}^\Omega(\psi))$, where $\Omega = (\omega^1, \omega^2, \omega^{12})$ with $\omega^1, \omega^2, \omega^{12} \subseteq \{1, \dots, n\}$, is equal to the corank of the matrix

$$\begin{bmatrix} S_{\omega^1} & \cdot \\ \cdot & S_{\omega^2} \\ S_{\omega^{12}} & S_{\omega^{12}} \end{bmatrix}, \quad (12)$$

and analogous results hold for arbitrary degrees. This shows that the invariants $\dim(\mathcal{V}_{n,r}^\Omega(\psi))$ can indeed be evaluated efficiently; moreover, the information needed to perform the calculation (i.e., the matrices S_ω) can easily be extracted from a generator matrix of the stabilizer.

There is in fact a close relationship between the LC invariants $\dim(\mathcal{V}_{n,r}^\Omega(\psi))$ and the polynomial LU invariants that we considered in [6]. In both cases, invariants correspond to dimensions of certain subspaces of $S_\psi \times \dots \times S_\psi$ that are defined in terms of constraints on the supports of their elements. In fact, if r equals 1 or 2 then the LC invariants in theorems 1 and 2 are also LU invariants; in other words, the invariants $\dim(\mathcal{V}_{n,r}^\Omega(\psi))$ and $|\mathcal{T}_{n,r}^\Omega(\psi)|$ take on equal values on LU equivalent stabilizer states. This property is well known (see e.g. [13]) for the invariants (8) and (11), corresponding to $r = 1$. In order to prove this assertion for $r = 2$, let $\Psi = |\psi\rangle\langle\psi|$ be (the projector associated with) a stabilizer state and take $\Omega = (\omega^1, \omega^2, \omega^{12})$. Then one can readily verify that $|\mathcal{V}_{n,2}^\Omega(\psi)|$ is, up to a multiplicative constant, equal to

$$\text{Tr} \{(\text{Tr}_{\bar{\omega}^1} \Psi) (\text{Tr}_{\bar{\omega}^2} \Psi) (\text{Tr}_{\bar{\omega}^{12}} \Psi)\}, \quad (13)$$

which is manifestly invariant under the action of local unitary operators (the operation Tr_ω denotes tracing out all qubits in the set ω). Moreover, this also shows that every $|\mathcal{T}_{n,2}^\Omega(\psi)|$ is an LU invariant, as it can be written as a linear combination with constant coefficients of the $|\mathcal{V}_{n,2}^\Omega(\psi)|$'s. Summarizing the current discussion, if r equals 1 or 2 then the invariants $\dim(\mathcal{V}_{n,r}^\Omega(\psi))$ and $|\mathcal{T}_{n,r}^\Omega(\psi)|$ take on equal values on LU equivalent stabilizer states. It is clear that this result is very interesting in relation with the problem whether LC equivalence and LU equivalence of stabilizer states are identical notions.

Finally, we wish to note that the two main results in this paper are to be regarded as *upper bounds* on the maximal r which needs to be considered in order to obtain a complete set of invariants. Indeed, it is likely that only invariants corresponding to (substantially) lower r need to be considered in order to recognize LC equivalence between two stabilizer states. One indication for

this conjecture is the proof of theorem 1, which shows that all the information of the LC equivalence class of a given state $|\psi\rangle$ is in fact contained within a single invariant $|\mathcal{T}_{n,n}^{\Omega*}(\psi)|$. A second state $|\psi'\rangle$ is LC equivalent to $|\psi\rangle$ if and only if

$$|\mathcal{T}_{n,n}^{\Omega*}(\psi)| = |\mathcal{T}_{n,n}^{\Omega*}(\psi')|, \quad (14)$$

(in fact, even $|\mathcal{T}_{n,n}^{\Omega*}(\psi')| > 0$ is sufficient) and the information in all other invariants is in fact redundant. It is not unlikely that there exist smaller complete lists of invariants (i.e. of smaller r) which exhibit less redundancies. Secondly, our work shows examples of LC equivalence classes which are indeed characterized by invariants of small r : e.g., any stabilizer state is LC equivalent to the GHZ state on n qubits if and only if the values of the invariants (11) coincide for both states [9]. We are therefore led to believe that the results in theorems 1 and 2 can in principle be improved - if not for all stabilizer states then at least for some interesting subclasses of states. It is in this context interesting to point out that it has been conjectured and later disproved in a graph theoretical context that the list of LC invariants (11) are sufficient to characterize the LC equivalence class of all stabilizer states [3, 14]. Therefore, invariants where r is at least 2 must be taken into account. It is to date not clear whether also higher r are to be considered.

Summarizing this paper, we have characterized LC equivalence classes of stabilizer states by means of a finite

set of invariants. Our invariants have simple descriptions within the binary stabilizer framework, in that they are equal to dimensions of certain subspaces of $S_\psi \times \dots \times S_\psi$. Furthermore, we discussed the link with existing families of LU invariants.

Acknowledgments

This research is supported by several funding agencies: Research Council KUL: GOA-Mefisto 666, GOA-Ambiorics, several PhD/postdoc and fellow grants; Flemish Government: - FWO: PhD/postdoc grants, projects, G.0240.99 (multilinear algebra), G.0407.02 (support vector machines), G.0197.02 (power islands), G.0141.03 (Identification and cryptography), G.0491.03 (control for intensive care glycemia), G.0120.03 (QIT), G.0452.04 (QC), G.0499.04 (robust SVM), research communities (ICCoS, ANMMM, MLDM); - AWI: Bil. Int. Collaboration Hungary/ Poland; - IWT: PhD Grants, GBOU (McKnow) Belgian Federal Government: Belgian Federal Science Policy Office: IUAP V-22 (Dynamical Systems and Control: Computation, Identification and Modelling, 2002-2006), PODO-II (CP/01/40: TMS and Sustainability); EU: FP5-Quprodis; ERNSI; Eureka 2063-IMPACT; Eureka 2419-FlITE; Contract Research/agreements: ISMC/IPCOS, Data4s, TML, Elia, LMS, IPCOS, Mastercard; QUIPROCON; QUPRODIS.

-
- [1] D. Gottesman. *Stabilizer codes and quantum error correction*. PhD thesis, Caltech, 1997. quant-ph/9705052.
 - [2] R. Raussendorf, D.E. Browne, and H.J. Briegel. Measurement-based quantum computation with cluster states. *Phys. Rev. A*, 68:022312, 2003. quant-ph/0301052.
 - [3] M. Hein, J. Eisert, and H.J. Briegel. Multi-party entanglement in graph states. *Phys. Rev. A*, 69:062311, 2004. quant-ph/0307130.
 - [4] W. Dür, H. Aschauer, and H.J. Briegel. Multiparticle entanglement purification for graph states. *Phys. Rev. Lett.*, 91:107903, 2003. quant-ph/0303087.
 - [5] M. Van den Nest, J. Dehaene, and B. De moor. Graphical description of the action of local clifford transformations on graph states. *Phys. Rev. A*, 69:022316, 2004. quant-ph/0308151.
 - [6] M. Van den Nest, J. Dehaene, and B. De Moor. Local invariants of stabilizer codes. *Phys. Rev. A*, 70:032323, 2004. quant-ph/0404106.
 - [7] M. Van den Nest, J. Dehaene, and B. De Moor. Efficient algorithm to recognize local clifford equivalence of graph states. *Phys. Rev. A*, 70:034302, 2004. quant-ph/0405023.
 - [8] F. Verstraete and J.I. Cirac. Valence bond solids for quantum computation. *Phys. Rev. A*, 70:060302, 2004. quant-ph/0311130.
 - [9] M. Van den Nest, J. Dehaene, and B. De moor. Local unitary versus local clifford equivalence of stabilizer states. *Phys. Rev. A*, 71:062323, 2005. quant-ph/0411115.
 - [10] E. Rains. Quantum codes of minimum distance two. *IEEE Trans. Inform. Theory*, 45(1):266–271, 1999. quant-ph/9704043.
 - [11] I. Chuang and M. Nielsen. *Quantum computation and quantum information*. Cambridge University press, Cambridge, 2000.
 - [12] P. Shor and R. Laflamme. Quantum analog of the macwilliams identities for classical coding theory. *Phys. Rev. Lett.*, 78(8):1600–1602, 1997. quant-ph/9610040.
 - [13] E.M. Rains. Quantum weight enumerators. *IEEE Trans. Inform. Theory*, 44(4):1388–1394, 1998. quant-ph/9612015.
 - [14] A. Bouchet. Recognizing locally equivalent graphs. *Discrete Math.*, 114(1-3):75–86, 1993.